


» Several studies have shown that glues have a reduced risk of early postoperative pain compared to traditional fixation methods (1-3).

- z Comparable recurrence rate than with traditional fixation methods (1-3).
- z Stand-alone fixation method in different hernia repair techniques (1-3).
- z Atraumatic fixation method reported to be better tolerated by the patient than traditional fixation methods (1-4).
- z Low intra- and postoperative morbidity (2, 3).
- z Decreased surgery time compared to traditional fixation methods (1, 3).
- z Good biocompatibility and in vivo tolerance (1-3, 5).
- z Very low rate of infections related to Histoacryl® (1-3, 5).
- z Efficient polymerization (6).



LAPAROSCOPIC  
SURGERY

**Histoacryl® LapFix**  
ATRAUMATIC MESH FIXATION  
FOR LAPAROSCOPIC SURGERY

Distributed by:  
B. Braun Medical N.V./S.A. | Lambroekstraat 5b | 1831 Diegem | Belgium  
+32 (0)2 712 86 50 | [customercare.be@bbraun.com](mailto:customercare.be@bbraun.com) | [www.bbraun.be](http://www.bbraun.be)

## AESCULAP® – a B. Braun brand

Aesculap AG | Am Aesculap-Platz | 78532 Tuttlingen | Germany  
Phone +49 7461 95-0 | Fax +49 7461 95-2600 | [www.aesculap.com](http://www.aesculap.com)

Manufacturer acc. to MDD 93/42/EEC  
B. Braun Surgical, S.A. | Carretera de Terrassa, 121 | 08191 Rubí | Spain

The main product trademark "Aesculap" and the product trademarks "Histoacryl" and "Optilene" are registered trademarks of Aesculap AG.

Subject to technical changes. All rights reserved. This brochure may only be used for the exclusive purpose of obtaining information about our products. Reproduction in any form partial or otherwise is not permitted.



# Histoacryl® LapFix

## ATRAUMATIC MESH FIXATION FOR LAPAROSCOPIC SURGERY



### REFERENCES / STUDIES

- (1) Hoyuela C, Juvany M, Carvajal F, Veres A, Troyano D, Trias M, Martrat A, Ardid J, Obiols J, López-Cano M. Randomized clinical trial of mesh fixation with glue or sutures for Lichtenstein hernia repair. Br J Surg. 2017;104(6):688-94.
- (2) Helbling C, Schlumpf R. Sutureless Lichtenstein: first results of a prospective randomized clinical trial. Hernia. 2003;7(2):80-4.
- (3) Kim-Fuchs C, Angst E, Vorburger S, Helbling C, Candinas D, Schlumpf R. Prospective randomized trial comparing sutured with sutureless mesh fixation for Lichtenstein hernia repair: long-term results. Hernia. 2012;16(1):21-7.
- (4) Liew W, Wai YY, Kosai NR, Gendeh HS. Tackers versus glue mesh fixation: an objective assessment of postoperative acute and chronic pain using inflammatory markers. Hernia. 2017;21(4):549-54.
- (5) Kukleta JF, Freytag C, Weber M. Efficiency and safety of mesh fixation in laparoscopic inguinal hernia repair using n-butyl cyanoacrylate: long-term biocompatibility in over 1,300 mesh fixations. Hernia. 2012;16(2):153-62.
- (6) Report ECT / Dr. Gabriel Siedle / Determination of breaking load of glued meshes, May 2011. Based on this test, it was confirmed that Histoacryl was fully polymerised during a waiting time of 10 or 60 minutes at room temperature.

# Histoacryl® LapFix

## ORDERING INFORMATION

Description	Article No.	Contents
Histoacryl® Lap Mesh Fixation: 1 unit with 2 Histoacryl® ampoules	1052008	2 x 0.5 ml 1 x LapFix applicator
Histoacryl® Lap Mesh Fixation: 5 units with 5 Histoacryl® ampoules	1050165	5 x 0.5 ml 5 x LapFix applicator
Histoacryl® Blue	1050044	5 units
Optilene® Mesh 7.5 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)	1065030	5 units
Optilene® Mesh 10 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)	1065040	5 units
Optilene® Mesh 15 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)	1065080	5 units
Optilene® Mesh LP blue striped 7.5 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)	1964715	5 units
Optilene® Mesh LP blue striped 10 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)	1964725	5 units
Optilene® Mesh LP blue striped 15 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)	1964705	5 units
Optilene® Mesh Elastic blue striped 10 x 15 cm (48 g/m <sup>2</sup> ; pore size 2.4 x 3.6 mm)	1964920	5 units

